

**A Note on Observations of Thermal Habitat Diversity  
at Channel Rehabilitation and Control Sites  
in the Trinity River During April 1998.**



Prepared by  
Sean P. Gallagher

U.S. Fish and Wildlife Service, Arcata Fish  
and Wildlife Office, Arcata, CA

A Note on Observations of Thermal Habitat Diversity  
at Channel Rehabilitation and Control Sites  
in the Trinity River During April 1998.

Prepared by

Sean P. Gallagher

U.S. Fish and Wildlife Service, Arcata Fish  
and Wildlife Office, Arcata, CA

April 1999

### Introduction

Habitat diversity, due to flow variation and geomorphic diversity, has been implicated as a major factor influencing bioproduction and biodiversity in river ecosystems (Poff et al. 1997, Richter et al. 1997, Richter et al. 1996a, Richter et al. 1996b, Stanford et al. 1996, Ward and Stanford 1996, Ligon et al. 1995). Water temperature, a macrohabitat factor, is generally considered for salmonids which require cold temperatures. However, thermal diversity may be a key factor for many other riverine species, especially as it relates to natural hydrological features. For instance, western pond turtles require slower moving water with higher temperatures (Reese and Welsh 1998). In the Trinity River an experimental approach to increasing habitat diversity has been employed which consists of widening the river by removal of the riparian encroached berm (USFWS 1997, 1994, Evans 1979). Nine sites in the river were rehabilitated between 1988 and 1993. Gallagher (1999), by comparing four of these sites to contemporaneous controls, found that habitat diversity (types and area), weighted usable area for some species, and density of some species and life stages was increased, with increasing flows, by channel rehabilitation. Water temperature and thermal habitat diversity have not been thoroughly examined in the Trinity River.

An aside to the study of channel rehabilitation, thermal diversity, not reported by Gallagher (1999) is reported here. This work was intended as a preliminary study and was conducted because temperature differences were discovered quite by accident during the course of the investigation of fish use and habitat diversity at rehabilitation sites (Gallagher 1999). A major hypothesis under consideration for this study was that channel rehabilitation increased habitat diversity. Micro-habitat was the major focus of the study (Gallagher 1999). However, thermal habitat diversity may be very important to river rehabilitation. This report presents preliminary findings supporting this hypothesis.

### Methods and Materials

Gallagher (1999), USFWS (1997), and McBain and Trush (1997) provide detailed descriptions of the study area and the study sites. During the week of 27 April 1998 water temperatures were measured at various locations in the main channel, along the bank, and in off channel areas at four restoration and control sites in the Trinity River California using a YSI temperature and conductivity meter. A data point recorded by Gallagher (1999) as part of an electro-fishing study of fish habitat use in the river was water conductivity. During the week of 27 April 1998, higher water temperatures were noted in an area of the Douglas City channel rehabilitation site. This spurred a quick preliminary check of water temperature along the banks of the rehabilitation and control sites during that week. Two persons, using a YSI temperature probe, worked up the bank at each site measuring water temperatures at a depth of approximately 0.06m in back waters, side channels, off channels and in the main channel. All temperatures were recorded in the afternoon between 1200 and 1700 hrs. The Lewiston Dam release during this week was 84.3 m<sup>3</sup>/s and the Douglas City gage reading was 94.6 m<sup>3</sup>/s. Portions of the riparian berm at control sites were inundated.

## Results

Air temperatures in the afternoon ranged from 19 to 26 C. At the Buck tail site the water temperature in the main channel was 9.5 C at both the rehabilitation and control sites. Water temperature along the constructed bank of the rehabilitation ranged from 10 to 18 C. Water temperatures downstream along the same bank at the control sites ranged from 11 to 13 C in the flooded riparian zone. Chinook salmon fry and juveniles, other fish species, and yellow legged frogs were observed in the warmer back water areas formed by channel rehabilitation at the Buck tail site. Figure 1 shows a sketch of the Buck tail site on 27 April 1998. This information was not collected systematically and statistical comparisons are not possible.

Water temperature differences were not as dramatic along the right bank (constructed bank at rehabilitation site) at the Lime Kiln site on 29 April 1998. Main channel water temperature was approximately 10 C. There was a narrow band of 11 to 11.5 C water along the constructed bank at this site. A small area of 12.5 C water was observed at the lower end of the site. Water temperature in the main channel at the Lime Kiln control site was also 10 C. In the back water areas formed by flooding of the riparian berm, water temperatures ranged from 11 to 18 C. In deeper areas thermal stratification was observed. Water at the surface was 18 C and along the bottom it was 13 C. Fish were observed in the warmer thermally stratified areas along the left bank at the Lime Kiln control site. Figure 2 is a sketch of the Lime Kiln rehabilitation and control sites on 29 April 1998.

Water temperature in the main channel at the Douglas City rehabilitation site and just down stream at a non-rehabilitated site was 11 C on 27 April 1998 at 1700 hrs. The rehabilitation site had a large backwater area connected at the top to the main channel by a small side channel. Water temperature in this area ranged from 15 to 21 C. Fish (chinook salmon young-of-the-year) were observed in this back water area. Water flowing at the top of the side channel feeding this backwater was 13 C. There were two other backwater areas formed by cobble bars that had temperatures higher than observed in the main channel. These measurements were 12.5 and 15 C. Downstream of the rehabilitation site in slow water areas along the bank water temperatures were the same as in the main channel (11 C). These areas were associated with the riparian berm and therefore experienced some shading during the day. The types of habitats created by water flowing over the rehabilitation site for this river discharge at the Douglas City rehabilitation site were similar to the Buck Tail site (Fig 1).

Water temperature in the main channel at the Bell Gulch rehabilitation and control sites was between 9.1 and 9.5 C on 30 April 1998 at 1200 hrs. In low velocity areas along the constructed bank at the rehabilitation site water temperatures were between 10 and 11 C. At the control site water temperatures in low velocity areas along the bank were the same as in the main channel. Physical habitat diversity at the Bell Gulch site was similar to the Lime Kiln site (Figs. 2, 3).

## Discussion

During the spring, the Trinity River historically experienced high, long duration snow melt flows which inundated cobble bars, riparian areas, and flooded backwaters and off channel areas. These areas probably supplied a mosaic of habitats with different thermal characteristics. The observations presented here support the notion that channel rehabilitation, as employed in the Trinity River, increases thermal habitat diversity by creating low velocity areas which are located off the main channel and experience differential heating. Thus habitat heterogeneity (Ward and Stanford 1995) is further increased. The temperature measurements reported here and the observations of fish (chinook salmon fry, brown trout fry, and stickle backs) and yellow legged frog use of the thermally stratified areas also support this idea. Murphy and Meehan (1991) suggest that removal of riparian vegetation can have immediate upstream benefits by increasing algae based food production but may limit overall production by cumulatively increasing temperature downstream. This is probably not of particular concern at this time as only a very small portion of the riparian channel has been modified. In addition, water temperatures in the Trinity are strongly controlled by Trinity Dam releases. The findings reported here suggest that small

areas of thermally diverse areas are created by rehabilitation at specific sites due to alluvial channel characteristics and bar formation. The finding that main channel water temperatures were the same between rehabilitation and downstream areas suggests that rehabilitation sites are not having a strong effect on main channel temperatures. The thermally diverse habitats we observed potentially provide important habitat for turtles, frogs, native and nonnative fish, and invertebrates. This type of habitat is currently scarce in the Trinity River due to the channeling effect of riparian berms and the effect of Trinity Dam release on water temperature. Shallow water areas with warm temperatures increase growth for young fish while decreasing predation by larger fish (Ward and Stanford 1995). Chapman and Bjorn (1969) found that during warmer months young chinook salmon and steelhead shift to faster deeper water as they grow. They also provide evidence of obvious diurnal-nocturnal changes in location for these species. Our findings did not qualify temperatures over time. However, fish at the rehabilitation sites may move and use thermally stratified areas during warmer and/or cooler periods. Diurnal thermal fluctuation in habitats created by inundation of cobble areas of rehabilitation sites may increase temporal habitat heterogeneity in the Trinity River. Thermal heterogeneity provides a mosaic of areas for fish to select from based on their physiological requirements. Thermal diversity may also result in an increase in macro-invertebrate production and diversity. Further experimental evaluation of the temperature aspects of channel rehabilitation in the Trinity River is needed. Diurnal as well as seasonal thermal diversity should be examined at rehabilitation and control sites. Species use of thermally diverse areas and their thermal requirements should also be investigated.

#### Acknowledgments

Jack Williamson of the U.S. Fish and Wildlife Service, Arcata, CA Fish and Wildlife Office and Troy Branham of the Hoopa Tribal Fisheries Department assisted with field data collection. Jay Glase, Rick Quihillalt, and Paul Zedonis of the U.S. Fish and Wildlife Service, Arcata, CA Fish and Wildlife Office provided helpful comments on earlier drafts of this note.

#### References

- Chapman, D. W. and T. C. Bjorn. 1969. Distribution of salmonids in streams, with special reference to food and feeding, pgs. 153-176. In T.G. Northcote (ed.) Symposium on salmon and trout in streams. H. R. Macmillan lectures in Fisheries. University of British Columbia, Vancouver, B. C.
- Evans, J. F. 1979. Evaluation of riparian vegetation encroachment Trinity River, California. Trinity River fish and wildlife task force.
- Gallagher, S. P. 1999. Experimental comparisons of fish habitat and fish use between channel rehabilitation sites and the vegetation encroached channel of the Trinity River.
- Ligon, F. K., W. E. Dietrich, and W. J. Trush. 1995. Downstream ecological effects of dams, a geomorphic perspective. *Bioscience*, 45: 183-192.
- McBain, S. and W. J. Trush. 1997. Trinity River maintenance flow study final report. McBain and Trush, Arcata, CA. 316 pp.
- Murphy, M. L. and W. R. Meehan. 1991. Influences of forest and range land management on salmonid fisheries and their habitats. *American Fisheries Society Special Publication* 19: 17-46.
- Poff, N. L., J. D. Allan, M. B. Bain, J. R. Karr, K. L. Prestegard, B. D. Richter, R. E. Sparks, and J. C. Stromberg. 1997. The natural flow regime a paradigm for river conservation and restoration. *Bioscience*, 47:769-784.
- Reese, D. A. and H. H. Welsh. 1998. Habitat use by western pond turtles in the Trinity River, California. *Journal of Wildlife Management*. 62: 842-853.
- Richter, B. D., J. V. Baumgartner, R. Wigington, and D. P. Braun. 1997. How much water does a river need. *Freshwater Biology*, 37:231-249.
- Richter, B. D., J. V. Baumgartner, J. Powell, and D. P. Braun. 1996a. A method for assessing hydrologic alteration within ecosystems. *Conservation Biology*, 10: 1163-1174.
- Richter, B. D., C. A. Frissell, R. N. Williams, J. A. Lichatowich, and C. C. Coutant. 1996b. A general protocol for restoration of regulated rivers. *Regulated Rivers: Research and Management*, 12: 391-413.

- Stanford, J. A., J. V. Ward, W. J. Liss, C. A. Frissell, R. N. Williams, J. A. Lichatowich, and C. C. Coutant. 1996. A general protocol for restoration of regulated rivers. *Regulated Rivers: Research and Management*, 12: 391-413.
- U.S.F.W.S. 1994. Restoration of the Mainstem Trinity River background report. Trinity River restoration program, US Fish and Wildlife Service. Weaverville, CA. 14 pp.
- U.S.F.W.S. 1997. Physical habitat and fish use of channel rehabilitation projects on the Trinity River. USFWS, Coastal California Fish and Wildlife Office, Arcata, CA, 19 pp.
- Ward, J. V. and J. A. Stanford. 1995. Ecological connectivity in alluvial river ecosystems and its disruption by flow regulation. *Regulated Rivers: Research and Management*. 11:105-119.

L1-28-418  
 Notes: TB  
 Temp: 5w  
 cfs: 2700  
 Air Ten 76°F

main flow

①  
Bridget  
②  
Tiers

1100  
@1527  
1300

East  
Tree

Back to 1  
control  
3144+  
Bact  
Pol in

Bucktail  
Feathered Edge

Handwritten note:  $14450 \pm 50$   
9.50 @ 145k

1381

11/5

1000 @ 1.15 =

1788  
1859

1704  
②  
1453

1202  
C  
1452

Yrsls

24

AND

AND

Land Trees

170C  
1441

1944	1945
1946	1947
1948	1949
1950	1951
1952	1953
1954	1955
1956	1957
1958	1959
1960	1961
1962	1963
1964	1965
1966	1967
1968	1969
1970	1971
1972	1973
1974	1975
1976	1977
1978	1979
1980	1981
1982	1983
1984	1985
1986	1987
1988	1989
1990	1991
1992	1993
1994	1995
1996	1997
1998	1999
2000	2001
2002	2003
2004	2005
2006	2007
2008	2009
2010	2011
2012	2013
2014	2015
2016	2017
2018	2019
2020	2021
2022	2023
2024	2025
2026	2027
2028	2029
2030	2031
2032	2033
2034	2035
2036	2037
2038	2039
2040	2041
2042	2043
2044	2045
2046	2047
2048	2049
2050	2051
2052	2053
2054	2055
2056	2057
2058	2059
2060	2061
2062	2063
2064	2065
2066	2067
2068	2069
2070	2071
2072	2073
2074	2075
2076	2077
2078	2079
2080	2081
2082	2083
2084	2085
2086	2087
2088	2089
2090	2091
2092	2093
2094	2095
2096	2097
2098	2099
2100	2101
2102	2103
2104	2105
2106	2107
2108	2109
2110	2111
2112	2113
2114	2115
2116	2117
2118	2119
2120	2121
2122	2123
2124	2125
2126	2127
2128	2129
2130	2131
2132	2133
2134	2135
2136	2137
2138	2139
2140	2141
2142	2143
2144	2145
2146	2147
2148	2149
2150	2151
2152	2153
2154	2155
2156	2157
2158	2159
2160	2161
2162	2163
2164	2165
2166	2167
2168	2169
2170	2171
2172	2173
2174	2175
2176	2177
2178	2179
2180	2181
2182	2183
2184	2185
2186	2187
2188	2189
2190	2191
2192	2193
2194	2195
2196	2197
2198	2199
2200	2201
2202	2203
2204	2205
2206	2207
2208	2209
2210	2211
2212	2213
2214	2215
2216	2217
2218	2219
2220	2221
2222	2223
2224	2225
2226	2227
2228	2229
2230	2231
2232	2233
2234	2235
2236	2237
2238	2239
2240	2241
2242	2243
2244	2245
2246	

10/15  
54  
54

Dr. [illegible]

16c  
⑤  
1215

1496  
P. 11105:60  
P. 11105:60

1602  
1446  
1701  
C

Handwritten notes:

- 130C
- 19°C surface
- 647
- 65J
- to 400

1 E110m  
Leagcd  
Frids

gravel

12

2





FIG #2



T. Bronham  
J. W. Williamson  
4/29/98  
2700 Down G  
1430 o'clock  
→ Temperature @ 0.2 depth

Bel Gulch FE H<sub>2</sub>O temp measure mnt  
Air temp ~ 72.

4/30/98 @ 1200

- 1 CPSE day temp 11°C depth 0.2 vel < 0.5
- 2 mcow 1 temp 9.5°C depth 0.5 vel < 0.5
- 3 CPSEW depth 0.2 temp 10°C vels < 0.2
- 4 mcow 2 temp 9.5°C depth 0.4 vels 1-1.5
- 5 CPSEW temp 10°C depth 0.2 waves

- 6 mcow 3 temp 9.9°C depth 0.3
- 7 CPSEW temp 10°C depth 0.2 waves

- 8 mcow 2 temp 9.5°C depth 0.4 vel ~ 1.5





Fig #3

Bel Gulch control

- 0 CPSEW temp 9.5°C depth 0.2 waves
- 2 mcow 1 temp 9°C depth 0.5
- 3 CPSEW temp 9.5°C depth 0.2 vels < 0.2
- 4 mcow 2 depth 0.4 temp 9.1°C vels ~ 0.5
- 5 CPSEW temp 9.5°C depth 0.2 vels < 0.1





# Appendix


340

#1

1650 hrs 4/27/98 DGE

H<sub>2</sub>O temp main channel ~ 0.8 f/s  
= 11°C measured ~ 10' out into current depth ~ 0.4

12 Ditch  
MSC



BACK H<sub>2</sub>O ~~off~~ connected to channel at top  
Through a flowing MSC H<sub>2</sub>O temp here is  
15°C. This is on near River side of  
BACK H<sub>2</sub>O CPBW. no flow here depth ~ 0.3

near BANK edge of same BACK H<sub>2</sub>O water temp is  
21°C! This is just ds of where MSC flows into  
This CPBW X's mark location of temp measurements!  
Low to zero flow here. Depth 0.2

In CPBW fish observed

y1 = top of MSC H<sub>2</sub>O temp 13°C flow ~ 0.2 f/s

y2 = 10' out in main channel H<sub>2</sub>O temp 11°C flow < 0.5 f/s  
This Area Between X7 & 3 US of X6 DS of main channel

main flow  
CPBW US of Xsec #5 H<sub>2</sub>O temp = 12.5°C no flow waves  
unit is 7 x 7 ft. main flow H<sub>2</sub>O temp = 10.5°C flow < 0.5 f/s depth 0.4 ft

CPBW just DS of Xsec #2 H<sub>2</sub>O temp in CPBW = 15°C waves  
but no flow unit is 12 x 20 ft. depth 0.2 ft

x main channel H<sub>2</sub>O temp = 11°C taken ~ 10 ft into channel  
~ 0.5 f/s flow depth ~ 0.4 ft

DS of FE @ flow study Xsec 3 Lg FEW no flow waves  
Depth 0.2 f 11°C H<sub>2</sub>O temp

DS of FE flow study Xsec #5 BAK FEW 0 flow waves  
H<sub>2</sub>O temp = 11°C Depth 0.2 ft.

BAK CPBW DS flow study X2 near old camp spot formed with MSC  
Temp measured with VSI 33 S-C-T meter